

# **MARINE MICROBIOLOGY**

## **ECOLOGY AND APPLICATIONS**

Second Edition

**Colin Munn**

With Foreword by Farooq Azam

# Contents

<b>Chapter 1 Microbes in the Marine Environment</b>	<b>1</b>	<b>Chapter 2 Methods in Marine Microbiology</b>	<b>25</b>
Marine microbiology is one of the most exciting and important areas of modern science	2	SAMPLING, GENERAL EXPERIMENTAL PROCEDURES, AND REMOTE SENSING	26
Marine microbiology encompasses all microscopic organisms and viruses	2	The aim of microbial ecology is the study of the diversity and activities of microbes <i>in situ</i>	26
Marine microbes are found in all three domains of cellular life	3	Measurement of specific cell constituents may be used as biomarkers of microbial activity	26
Horizontal gene transfer confounds our understanding of evolution	4	Remote sensing and sampling permits analysis of microbial activities	27
Viruses are noncellular entities with great importance in marine ecosystems	4	Microbiological sampling requires special techniques	27
Microbial processes shape the living world	5	Mecocosm experiments attempt to simulate natural conditions	29
Marine microbes show great variation in size	5	Microelectrodes and biosensors are used to measure environmental changes	30
The world's oceans and seas form an interconnected water system	7	Isotopes are used to study microbial transformations of compounds	30
The upper surface of the ocean is in constant motion owing to winds	9	DIRECT OBSERVATION AND ENUMERATION OF MICROBES	31
Deep-water circulation systems transport water between the ocean basins	10	Light and electron microscopy are used to study morphology and structure of microbes	31
Seawater is a complex mixture of inorganic and organic compounds	10	Epifluorescence light microscopy enables enumeration of marine microbes	32
Light and temperature have important effects on microbial processes	11	Confocal laser scanning microscopy enables recognition of living microbes within their habitat	33
Marine microbes form a major component of the plankton	13	Flow cytometry measures the number and size of particles	33
Microbes, particles, and dissolved nutrients are not evenly distributed in seawater	14	CULTURE-BASED METHODS FOR ISOLATION AND IDENTIFICATION OF MICROBES	35
Microbes play a key role in the formation of sediments	16	Different microbes require specific culture media and conditions for growth	35
Microbes colonize surfaces through formation of biofilms	17	Enrichment culture selects for microbes with specific growth requirements	36
Microbes in sea ice form an important part of the food chain in polar regions	19	Phenotypic testing is used for identification and detailed characterization of many cultured bacteria	38
Microbial activity at hydrothermal vents provides an oasis of life in the deep sea	19	Analysis of microbial components can be used for bacterial classification and identification	38
Cold seeps also support diverse life	20		
Living organisms are the habitats of many microbes	21		
Conclusions	22		
References	22		
Further reading	23		

NUCLEIC-ACID-BASED METHODS	40	Anaerobic anoxygenic photosynthesis uses only one type of reaction center	62
The use of nucleic-acid based methods has had a major impact on the study of marine microbial diversity	40	Aerobic anoxygenic phototrophy is widespread in planktonic bacteria	62
Sequencing of ribosomal RNA genes is the most widely used tool in studies of microbial diversity	40	Some phototrophs use rhodopsins as light-harvesting pigments	63
The first step in all nucleic acid investigations involves the isolation of genomic DNA or RNA from the culture or community	42	Chemolithotrophs use inorganic electron donors as a source of energy and reducing power	63
The polymerase chain reaction (PCR)	42	Thiotrophic bacteria use sulfur compounds as electron donor	63
Genomic fingerprinting is used for detailed analysis of cultured microbes	44	Many chemolithotrophs use hydrogen as an electron donor	65
Determination of GC ratios and DNA–DNA hybridization is used in bacterial taxonomy	46	Nitrification by <i>Bacteria</i> and <i>Archaea</i> is a major process in the marine nitrogen cycle	65
DNA sequencing is a major tool in marine microbiology	46	The Calvin–Benson cycle is the main method of carbon dioxide fixation in autotrophs	66
“Next-generation” technologies allow inexpensive high-throughput sequencing	46	Some <i>Archaea</i> and <i>Bacteria</i> use alternative pathways to fix CO <sub>2</sub>	66
Sequence data are used for phylogenetic analysis	47	Fixation of nitrogen makes this essential element available for building cellular material in all life	67
Denaturing gradient gel electrophoresis (DGGE) and terminal restriction fragment length polymorphism (TRFLP) are widely used to assess composition of microbial communities	48	Many marine microbes obtain energy by the fermentation of organic compounds	68
Elucidating the full genome sequence of microbes has provided major insights into their functional roles	49	Aerobic and anaerobic respiration use external electron acceptors	68
Metagenomics is revolutionizing our understanding of marine microbial ecology	51	Reduction of nitrate and denitrification result in release of nitrogen and other gases	69
Fluorescent hybridization (FISH) allows visualization and quantification of specific microbes	53	Sulfate reduction is a major process in marine sediments	69
Metatranscriptomics and metaproteomics reveal metabolic activities in the environment	54	Methanogenesis is a special type of metabolism carried out only by a group of <i>Archaea</i>	70
Microarrays enable assessment of gene activity in the environment	55	Aerobic catabolism of methane and other C <sub>1</sub> compounds is widespread in coastal and oceanic habitats	70
Conclusions	55	Use of complex macromolecules requires the synthesis of extracellular enzymes	71
References	55	Acquisition of iron is a major challenge for marine microbes	71
Further reading	56	The growth of bacterial cells depends on availability of nutrients and environmental factors	72
<b>Chapter 3 Metabolic Diversity and Ecophysiology</b>	<b>59</b>	Bacteria adapt to starvation by a series of coordinated changes to cell metabolism	72
All cells need to obtain energy and conserve it in the compound ATP	60	Most marine microbes are adapted to an oligotrophic lifestyle and grow very slowly	73
All cells need carbon as the major component of cellular material	60	Some bacteria enter a “viable but nonculturable” state in the environment	73
Phototrophy involves conversion of light energy to chemical energy	60	Nutrients are acquired via specialized transport mechanisms	74
Oxygenic photosynthesis involves two distinct but coupled photosystems	61	Growth efficiency of many marine bacteria is probably low	75

Microbes use a variety of mechanisms to regulate cellular activities	75	A wide range of <i>Proteobacteria</i> can grow chemolithotrophically using reduced sulfur compounds	101
Some bacteria use motility in the quest for nutrients and optimal conditions	76	Aerobic methanotrophs and methylotrophs are widespread in coastal and oceanic habitats	102
Formation of biofilms is an important step in microbial colonization of surfaces	79	The pseudomonads are a heterogeneous group of chemoorganotrophic, aerobic, rod-shaped <i>Proteobacteria</i>	103
Pili are important for bacterial attachment to surfaces and exchange of genetic information	80	Free-living aerobic nitrogen-fixing bacteria are important in sediments	103
Antagonistic interactions between microbes occur on particles or surfaces	80	The <i>Enterobacteriaceae</i> is a large and well-defined family of <i>Gammaproteobacteria</i>	103
Quorum sensing is an intercellular communication system for regulation of gene expression	81	<i>Vibrio</i> and related genera have worldwide distribution in coastal and ocean water and sediments	104
Most marine microbes grow at low temperatures	82	Some members of the <i>Vibrionaceae</i> are bioluminescent	104
Microbes growing in hydrothermal systems are adapted to very high temperatures	84	The <i>Oceanospiralles</i> are characterized by their ability to break down complex organic compounds	106
Microbes that inhabit the deep ocean must withstand a very high hydrostatic pressure	85	Magnetotactic bacteria orient themselves in the Earth's magnetic field	107
Microbes vary in their requirements for oxygen or tolerance of its presence	86	<i>Bdellovibrio</i> is a predator of other bacteria	108
Ultraviolet irradiation has lethal and mutagenic effects	87	Budding and stalked <i>Proteobacteria</i> show asymmetric cell division	108
Microbes are protected from osmotic damage by various mechanisms	87	Sulfur- and sulfate-reducing bacteria have a major role in the sulfur cycle	109
Conclusions	88	The <i>Cyanobacteria</i> carry out oxygenic photosynthesis	110
References	88	Many marine <i>Cyanobacteria</i> carry out nitrogen fixation	111
Further reading	89	The genera <i>Prochlorococcus</i> and <i>Synechococcus</i> dominate the picoplankton in large areas of the Earth's oceans	112
<b>Chapter 4 Marine Bacteria</b>	<b>91</b>	<i>Cyanobacteria</i> are important in the formation of microbial mats in shallow water	114
OVERVIEW OF DIVERSITY OF THE BACTERIA	92	The <i>Firmicutes</i> are a major branch of Gram-positive <i>Bacteria</i>	114
The domain <i>Bacteria</i> contains about 80 phyla, many of which have no cultivated members	92	<i>Epulopiscium fishelsoni</i> and related species are giant bacteria with a unique "viviparous" lifestyle	115
There is no generally accepted concept for the definition of bacterial species	93	The <i>Actinobacteria</i> is a large phylum including the mycobacteria and actinomycetes	116
<i>Bacteria</i> show a variety of cell forms and structure	93	The <i>Cytophaga-Flavobacterium-Bacteroides</i> group is morphologically and metabolically diverse	116
The cell wall is an important feature of bacterial cells	94	The <i>Planctomycetes</i> are a group with cells that show some similarities to eukaryotes	117
Many <i>Bacteria</i> produce a glycocalyx or capsule	96	<i>Verrucomicrobia</i> is a poorly characterized phylum of <i>Bacteria</i>	118
Phylogenetic studies of planktonic <i>Bacteria</i> reveal a small number of major clades	96		
MAJOR TYPES OF MARINE BACTERIA, GROUPED BY PHENOTYPE	98		
Several groups of bacteria carry out anoxygenic photosynthesis	98		
Nitrifying bacteria grow chemolithotrophically using reduced inorganic nitrogen compounds as electron donors	99		

The spirochetes are Gram-negative, tightly coiled, flexuous bacteria distinguished by very active motility	118	The choanoflagellates have a unique feeding mechanism	138
<i>Aquifex</i> and <i>Thermotoga</i> are hyperthermophiles	118	Dinoflagellates have critical roles in marine systems	138
Conclusions	119	Dinoflagellates undertake diurnal vertical migration	139
References	119	Some dinoflagellates exhibit bioluminescence	139
Further reading	120	The ciliates are voracious grazers of other protists and bacteria	140
<b>Chapter 5 Marine Archaea</b>	<b>121</b>	The haptophytes (prymnesiophytes) are major components of ocean phytoplankton	141
Several aspects of cell structure and function distinguish the <i>Archaea</i> from the <i>Bacteria</i>	122	Diatoms are extremely diverse and abundant primary producers in the oceans	144
The <i>Euryarchaeota</i> and <i>Crenarchaeota</i> form the major branches of the <i>Archaea</i>	122	Diatoms and their products—past and present—have many applications	145
Many members of the <i>Euryarchaeota</i> produce methane	122	Protists in the picoplankton size range are extremely widespread and diverse	145
<i>Archaea</i> in deep sediments can carry out anaerobic oxidation of methane coupled to sulfate reduction	124	Raphidophytes are stramenopiles which may cause harmful blooms	146
<i>Thermococcus</i> and <i>Pyrococcus</i> are hyperthermophiles found at hydrothermal vents	125	Thraustochytrids and labyrinthulids play an important role in breakdown and absorption of organic matter	147
<i>Archaeoglobus</i> and <i>Ferroglobus</i> are hyperthermophilic sulfate-reducers and iron-oxidizers	126	Amoebozoa may be important grazers of bacteria associated with particles	148
Some <i>Euryarchaeota</i> exist in hypersaline environments	126	Radiolarians and foraminifera have highly diverse morphologies with mineral shells	148
<i>Nanoarchaeum</i> is an obligate parasite of another archaeon, <i>Ignicoccus</i>	127	Marine fungi are especially important in decomposition of complex materials in coastal habitats	149
The <i>Crenarchaeota</i> include hyperthermophiles and psychrophiles	127	Conclusions	151
Hyperthermophilic <i>Crenarchaeota</i> belong to the order <i>Desulfurococcales</i>	127	References	151
The psychrophilic marine <i>Crenarchaeota</i> are major members of the plankton	128	Further reading	152
Conclusions	129	<b>Chapter 7 Marine Viruses</b>	<b>153</b>
References	131	Viruses are extremely diverse in structure and genetic composition	154
Further reading	131	Viruses are the most abundant biological entities in seawater	155
<b>Chapter 6 Marine Eukaryotic Microbes</b>	<b>133</b>	Phages are viruses that infect bacterial and archaeal cells	157
The term “protist” is used to describe an extremely diverse collection of unicellular eukaryotic microbes	134	The life cycle of phages shows a number of distinct stages	158
Systems for the classification of eukaryotic microbes are still developing	135	Lysogeny occurs when the phage genome is integrated into the host genome	160
Many protists possess flagella	135	Large DNA viruses are important pathogens of planktonic protists	161
The euglenids may be phototrophic, heterotrophic, or mixotrophic	136	Photosynthetic protists are also infected by RNA viruses	163
The bicosoecids are a group of highly active bacterivorous flagellates	137	The role of viruses as pathogens of heterotrophic protists remains unclear	164

Loss of infectivity of viruses arises from irreparable damage to the nucleic acid or protein capsid	165	Ocean microbes require iron	189
Measurement of virus production rates is important for assessing the role of virus-induced mortality	165	<b>THE NITROGEN CYCLE</b>	190
Viral mortality “lubricates” the biological pump	166	Major shifts in our understanding of the marine nitrogen cycle are in progress	190
Viral mortality plays a major role in structuring diversity of microbial communities	166	New nitrogen-fixers have been discovered recently	191
Marine viruses show enormous genetic diversity	167	Fixed nitrogen is returned to the inorganic pool by ammonification and nitrification	191
Viromes are reservoirs of genetic diversity and exchange	167	Denitrification and anammox reactions return nitrogen to its elemental form	192
Conclusions	168	Microbial processes in sediments are a major contributor to nitrogen cycling	193
References	170	<b>THE SULFUR CYCLE</b>	193
Further reading	171	The oceans contain large quantities of sulphur—an essential element for life	193
<b>Chapter 8 Microbes in Ocean Processes—Carbon Cycling</b>	<b>173</b>	Metabolism of organic sulfur compounds is especially important in surface waters	193
Development of the microbial loop concept transformed our understanding of ocean processes	174	A fraction of DMSP production leads to release of the gas dimethyl sulfide (DMS)	196
The fate of carbon dominates consideration of the microbial ecology of the oceans	175	Microbial sulfate reduction and sulfide oxidation occur in sediments, vents, and seeps	196
Marine phytoplankton are responsible for about half of the global CO <sub>2</sub> fixation	175	<b>THE PHOSPHORUS CYCLE</b>	197
As well as light, photosynthetic activity depends on the availability of nutrients	178	Phosphorus is often a limiting or colimiting nutrient	197
The importance of various components of the microbial loop varies according to circumstances	179	Marine microbes are adapted to low and variable levels of phosphorus	200
The microbial loop results in retention of dissolved nutrients	182	Conclusions	201
Ingestion of bacteria by protists plays a key role in the microbial loop	183	References	201
The “viral shunt” catalyzes nutrient regeneration in the upper ocean	184	Further reading	202
Eutrophication of coastal waters affects microbial activity	185	<b>Chapter 10 Symbiotic Associations</b>	<b>203</b>
Conclusions	185	Zooxanthellae and other photosynthetic endosymbionts are vital for the nutrition of many marine animals	204
References	186	Coral bleaching occurs due to the breakdown of the symbiosis between zooxanthellae and their host	205
Further reading	186	Scleractinian corals are multipartner symbiotic systems (holobionts)	206
<b>Chapter 9 Microbes in Ocean Processes—Nitrogen, Sulfur, Iron, and Phosphorus Cycling</b>	<b>187</b>	Photosynthetic zooxanthellae boost the growth of giant clams in nutrient-poor waters	207
<b>NUTRIENT LIMITATION</b>	188	Worms and clams at hydrothermal vents obtain nutrition from chemosynthetic bacterial endosymbionts	207
Key elements may act as limiting nutrients for different groups of microbes	188	Chemosynthetic symbionts are widely distributed in marine invertebrates	209
Productivity of surface waters shows marked geographical variations	188	Animals colonizing whale falls depend on autotrophic and heterotrophic symbionts	212

Some hydrothermal vent animals have dense populations of bacteria on their surface	212	Parasitic dinoflagellates are major pathogens of crustaceans	235
Some fish and invertebrates use bacteria to make light	213	<b>DISEASES OF VERTEBRATES</b>	<b>235</b>
The bobtail squid uses bacterial bioluminescence for camouflage	214	Microbial diseases of fish cause major losses in aquaculture, but effects on natural populations are harder to determine	235
Endosymbionts of bryozoans produce compounds that protect the host from predation	215	The importance of fish diseases in aquaculture has led to the development of specialized branches of veterinary science and diagnostic microbiology	236
Sponges contain dense communities of specific microbes	215	Bacteria produce infections in fish using a range of pathogenic mechanisms	238
Some protists with endosymbionts can switch from heterotrophic to phototrophic metabolism	218	Vibrios are responsible for some of the main infections of marine fish	238
Viruses may help a sea slug to use “stolen” chloroplasts for photosynthesis	218	Pasteurellosis is a major disease in warm-water marine fish	241
Conclusions	219	<i>Aeromonas salmonicida</i> has a broad geographic range affecting fish in fresh and marine waters	241
References	219	Marine flexibacteriosis is caused by an opportunist pathogen of low virulence	243
Further reading	220	<i>Piscirickettsia</i> and <i>Francisella</i> are intracellular proteobacteria causing economically important diseases in salmon and cod	243
<b>Chapter 11 Microbial Diseases of Marine Organisms</b>	<b>223</b>	Intracellular Gram-positive bacteria cause chronic infections of fish	244
<b>DISEASES OF INVERTEBRATES</b>	<b>224</b>	Several Gram-positive cocci cause diseases affecting the central nervous system of fish	245
Diseases of invertebrates have major ecological and economic impact	224	Viruses cause numerous diseases of marine fish	245
Infectious diseases of corals have emerged as a major threat to their survival	224	Infectious salmon anemia virus is one of the most important pathogens in salmon culture	246
The fungus <i>Aspergillus sydowii</i> caused a mass mortality of sea fans in the Caribbean Sea	226	Viral hemorrhagic septicemia virus infects many species of wild fish	246
Black band disease of corals is a long-established disease of corals worldwide	226	Lymphocystis virus causes a highly contagious chronic skin infection of fish	247
White plague and white pox are major diseases affecting Caribbean reefs	227	Birnaviruses appear to be widespread in marine fish and invertebrates	247
Extensive tissue necrosis of corals may involve bacteria and protistan parasites	227	Viral nervous necrosis is an emerging disease with major impact	247
The role of viruses in coral diseases is unclear	227	Protists can cause disease in fish via infections, toxins, and direct physical effects	247
Sponge disease is a poorly investigated global phenomenon	228	Dinoflagellate and diatom toxins can affect marine mammals	248
Vibrios are a major cause of important diseases of cultured mollusks	228	Mass mortalities in the late twentieth century prompted the study of viral diseases of marine mammals	249
A wide range of other bacteria can cause infections in bivalve mollusks	229	Viruses from nine different families have been linked to diseases of marine mammals	250
Virus infections are a major problem in oyster culture	232	Several species of bacteria and fungi infect marine mammals	251
Bacterial and viral diseases are major problems in aquaculture of crustaceans	232	Sea turtles are affected by a virus promoting growth of tumors	251
Expansion of intensive prawn culture has been accompanied by a dramatic spread in viral diseases	233		
Bacteria can cause epizootics with high mortalities in crustaceans	234		

DISEASES OF SEaweEDS AND SEagrASSES	252	Brevetoxin can cause illness via ingestion or inhalation during red tides	272
Fungi, bacteria, and protists cause ecologically and economically important diseases of seaweeds and seagrasses	252	Diarrhetic shellfish poisoning and azaspiracid poisoning result in gastrointestinal symptoms	273
Many species of algae contain virus-like particles	253	Amnesic shellfish poisoning is caused by toxic diatoms	274
Conclusions	254	Ciguatera fish poisoning has a major impact on the health of tropical islanders	274
References	254	Dinoflagellates and diatoms probably produce toxins as antipredator defense mechanisms	275
Further reading	256	The incidence of harmful algal blooms and toxin-associated diseases is increasing owing to the interaction of many complex factors	
<b>Chapter 12 Marine Microbes as Agents of Human Disease</b>	<b>259</b>	Coastal waters must be regularly monitored to assess the development of harmful algal blooms	277
Pathogenic vibrios are common in marine and estuarine environments	260	Conclusions	279
Cholera is a major human disease with a reservoir in coastal environments	260	References	279
<i>Vibrio cholerae</i> produces disease in humans owing to production of a toxin and other pathogenic factors	261	Further reading	281
Control of cholera remains a major world health problem	261	<b>Chapter 13 Microbial Aspects of Marine Biofouling, Biodeterioration, and Pollution</b>	<b>283</b>
Mobile genetic elements play a major role in the virulence of <i>Vibrio cholerae</i>	262	BIOFOULING AND BIODETERIORATION	284
Non-O1 and non-O139 serotypes of <i>Vibrio cholerae</i> are widely distributed in coastal and estuarine waters	263	Microbial biofilms are often the first phase in biofouling	284
<i>Vibrio vulnificus</i> causes serious illness associated with seafood	263	Microbially induced corrosion occurs as a result of the activities of microorganisms within biofilms on metals, alloys and composite materials	284
Distribution of <i>Vibrio vulnificus</i> in the marine environment is affected by temperature and salinity	265	Microbes cause biodeterioration of marine wooden structures and timber	285
<i>Vibrio vulnificus</i> and other marine vibrios can cause wound infections	265	Microbial growth and metabolism are the major cause of spoilage of seafood products	285
Seafood-borne infection by <i>Vibrio parahaemolyticus</i> is common throughout the world	266	Processing, packaging, and inhibitors of spoilage are used to extend shelf-life	286
Scombroid fish poisoning is a result of bacterial enzymic activity	267	Some seafood products are produced by deliberate manipulation of microbial activities	287
Botulism is a rare lethal intoxication from seafood	268	MICROBIAL ASPECTS OF MARINE POLLUTION BY SEWAGE	287
Fugu poisoning is caused by a neurotoxin of probable bacterial origin	268	Coastal pollution by wastewater is a significant source of human disease	287
Some diseases of marine mammals and fish can be transmitted to humans	269	A range of human viruses are present in seawater contaminated by sewage	288
Toxic dinoflagellates and diatoms pose serious threats to human health	270	Fecal indicator bacteria have been used for many years to test public health risks in marine water	289
Paralytic shellfish poisoning is caused by saxitoxins produced by dinoflagellates	270	<i>Escherichia coli</i> and coliforms are unreliable indicators of human fecal pollution of the sea	291
Management of paralytic shellfish poisoning depends on assaying toxins in shellfish	271		



The fecal streptococci or enterococci are more reliable indicators for monitoring marine water quality	291	Metagenomics and bioinformatics lead to new biotechnological developments	311
Molecular-based methods permit quicker analysis of indicator organisms and microbial source tracking	292	Polymers from marine bacteria are finding increasing applications	312
A variety of alternative indicator species have been investigated	293	Microalgae are promising new sources of biofuels	312
Different countries use different quality standards for marine waters	294	Marine microbes are a rich source of biomedical products	313
Sewage pollution of water in which shellfish are harvested for human consumption poses a serious health hazard	294	Bioactive compounds from marine invertebrates may be produced by microbial symbionts	313
Many countries have microbiological standards for the classification of waters in which shellfish are cultivated	297	New antimicrobial compounds may be discovered through study of complex microbial communities	315
Direct testing for pathogens in shellfish is possible using molecular techniques	297	Marine microbes are the source of a range of health-promoting products	315
<b>OIL AND OTHER CHEMICAL POLLUTION</b>	298	New approaches to antifouling have been discovered through study of microbial colonization of surfaces	316
Oil pollution of the marine environment is a major problem	298	Marine microbes are a rich source for biomimetics, nanotechnology, and bioelectronics	316
A range of microbes are responsible for biodegradation of oil at sea	299	Microbial biotechnology has many applications in aquaculture	317
The fate of oil depends on a combination of physical and biological processes	299	Most bacterial pathogens can be killed or inhibited by antimicrobial agents	319
Biodegradation is enhanced by addition of emulsifiers	301	Resistance to antimicrobial agents is a major problem in aquaculture	320
Addition of nutrients is necessary to increase the rate of oil biodegradation	301	Vaccination of finfish is widely used in aquaculture	321
Bioremediation has been used to lessen the impact of oil spills on vulnerable coasts	302	Recombinant DNA technology is used to produce vaccines for diseases caused by viruses and some bacteria	322
Microbes are important in the distribution of persistent organic pollutants	302	DNA vaccination or genetic immunization depends on expression of a sequence encoding the protective antigen	323
Bacteria are effective in the removal of heavy metals from contaminated sediments	303	Probiotics, prebiotics, and immunostimulants are widely used in marine aquaculture	326
Microbial systems can be used for monitoring the environment for toxic chemicals	303	Conclusions	327
Mobilization of mercury by bacterial metabolism leads to accumulation of toxic methylmercury	304	References	327
Conclusions	304	Further reading	329
References	305	<b>Chapter 15 Concluding Remarks</b>	<b>331</b>
Further reading	306	<b>Glossary</b>	<b>333</b>
<b>Chapter 14 Marine Microbes and Biotechnology</b>	<b>307</b>	<b>Abbreviations</b>	<b>341</b>
Enzymes from marine microbes have many applications	308	<b>Index</b>	<b>343</b>
DNA polymerases from hydrothermal vent organisms are widely used in molecular biology	310	<b>Color plates</b>	<b>367</b>