

Estimates of the Influence of Radionuclide Solubility Limits and Sorption Competition on the Sorption Values in the SDBs of MX-80 Bentonite and Opalinus Clay

**Nuclear Energy and Safety Research Department (NES)
Laboratory for Waste Management (LES)**

M.H. Bradbury, M. Marques Fernandes,
B. Baeyens

Paul Scherrer Institut
Forschungsstrasse 111
5232 Villigen PSI, Switzerland
Tel. +41 56 310 21 11
Fax +41 56 310 21 99
www.psi.ch

PSI Bericht Nr. 17-04
November 2017
ISSN 1019-0643

Table of Contents

Preface	I
Abstract	III
Zusammenfassung.....	V
Résumé	VII
Table of Contents	IX
List of Tables.....	X
List of Figures	XI
1 Introduction	1
2 Background	3
3 Modifications to the 2SPNE SC/CE model.....	5
4 Factors influencing the sorption of radionuclides in real systems	7
4.1 Presence of stable isotopes	7
4.2 Complexation with anionic ligands in real systems.....	7
4.3 Summary.....	10
5 Illustrative example of a repository near-field competitive sorption scenario	11
5.1 General.....	11
5.2 Maximum limit concentrations.....	11
5.3 Calculation procedure	12
5.4 Divalent metals	13
5.5 Trivalent metals	16
5.6 Tetravalent metals.....	18
5.6.1 Tc(IV), Po(IV) and Zr(IV).....	18
5.6.2 Sn(IV)	19
5.6.3 Th(IV), Np(IV) and U(IV).....	19
6 The influence of competitive sorption on sorption values in Opalinus Clay.....	21
6.1 Divalent metals	21
6.2 Trivalent metals	22
6.3 Tetravalent metals.....	23
7 Summary of results and conclusions	25
8 References.....	27

Appendix A: Metal hydrolysis and surface complexation data for montmorillonite and illite	A-1
Appendix B: Auxiliary thermodynamic data	B-1
Appendix C: MX-80 mineralogy and porewater chemistry	C-1
Appendix D: Solubility limits of safety-relevant elements in the SF/HLW bentonite near-field	D-1
Appendix E: Opalinus Clay mineralogy and porewater chemistry	E-1

List of Tables

Tab. 4.1: Porewater compositions at pH = 7.25.....	9
Tab. 4.2: Eu speciation in the simple background electrolyte (Tab. 4.1, Case A) and in a complex porewater composition at pH = 7.25 (Tab. 4.1, Case B).....	9
Tab. 5.1: "Maximum limit concentrations" of metals in the porewater of a bentonite near-field of a high level waste repository (Tab. D-1 in Appendix D).....	12
Tab. 5.2: Summary of calculations for divalent elements (Group 1, Tab. 5.1).....	14
Tab. 5.3: Summary of calculations for trivalent elements (Group 2, Tab. 5.1).	16
Tab. 5.4: Summary of calculations for tetravalent elements (Group 3, Tab. 5.1).....	20
Tab. 6.1: Ni(II) sorption in Opalinus Clay at trace concentration (no competition) and in the presence of 5.2×10^{-5} M Fe(II).....	22
Tab. 6.2: Comparison of sorption values in Opalinus Clay for trivalent actinides / lanthanides calculated at trace concentrations with no competition and at maximum limit concentrations with competition.	22
Tab. 6.3: Comparison of sorption values in Opalinus Clay for tetravalent actinides calculated at trace concentrations with no competition and at maximum limit concentrations with competition.	23
Tab. 7.1: Comparison of sorption values for trivalent lanthanides and actinides considering their maximal limit concentrations and taking into account competitive sorption.	26
Tab. 7.2: Comparison of sorption values for tetravalent actinides considering their maximal limit concentrations and taking into account competitive sorption.	26
Tab. A-1: Hydrolysis and surface complexation reactions and constants for Ni(II) and Fe(II).....	A-1
Tab. A-2: Hydrolysis and surface complexation reactions and constants for Eu/Sm/Ho(III), Am/Ac/Cm(III) and Pu(III).	A-1

Tab. A-3:	Hydrolysis and surface complexation reactions and constants for Th(IV), U(IV) and Np(IV).....	A-2
Tab. B-1:	Aqueous thermodynamic data for divalent metals taken from the PSI/Nagra TDB 12/07 (Thoenen et al. 2014).....	B-1
Tab. B-2:	Aqueous thermodynamic data for trivalent metals taken from the PSI/Nagra TDB 12/07 (Thoenen et al. 2014).....	B-1
Tab. B-3:	Aqueous thermodynamic data for tetravalent metals taken from the PSI/Nagra TDB 12/07 (Thoenen et al. 2014).....	B-2
Tab. C-1:	Reference MX-80 mineralogy (Müller-Vonmoos & Kahr 1983).....	C-1
Tab. C-2:	Bentonite porewater obtained after 10'000 years of exchange between the bentonite starting system and the Opalinus Clay reference system without a low-pH shotcrete liner.	C-1
Tab. D-1:	Solubility limits (in mol/kg H ₂ O) of safety-relevant elements in the SF/HLW near-field (bentonite).	D-1
Tab. E-1:	Reference mineralogy and values for the bounding minerals for Opalinus Clay (Traber & Blaser 2013).....	E-1
Tab. E-2:	Porewater composition of Opalinus Clay (Mäder 2009).....	E-1

List of Figures

Fig. 4.1:	Eu isotherms on montmorillonite in two different water compositions at pH = 7.25.	8
Fig. 5.1:	Fe(II) sorption isotherm and the sorption of trace Zn in the presence of increasing Fe(II) concentrations ($Z_{\text{tot}} = 3 \times 10^{-8}$ M) on a synthetic iron-free montmorillonite (Soltermann et al. 2014).	13
Fig. 5.2:	Calculated Ni sorption isotherm for MX-80 in the bentonite porewater (pH 7.79); (a), (b) in the absence of competing metals and (c), (d) in the presence of Fe(II) at siderite saturation, 3.7×10^{-5} M.	15
Fig. 5.3:	Calculated Eu sorption isotherm for MX-80 in the bentonite porewater (pH 7.79) (a), (b) in the absence of competing metals and (c), (d) in the presence of competing metals whose equilibrium concentrations are as given in Tab. 5.1.....	17
Fig. 5.4:	Aqueous speciation of Tc(IV) in 0.1 M NaClO ₄ as function of pH.....	18
Fig. 5.5:	Aqueous speciation of Zr(IV) in 0.1 M NaClO ₄ as function of pH.....	18
Fig. 5.6:	Aqueous speciation of Sn(IV) in 0.1 M NaClO ₄ as function of pH.	19
Fig. 5.7:	Aqueous speciation of Th(IV), U(IV) and Np(IV) in 0.1 M NaClO ₄ as a function of pH.....	20
Fig. 6.1:	Calculated Ni sorption isotherm in the Opalinus Clay / porewater (pH 7.20) system.	21