

Diffusion of Selected Cations and Anions in Compacted Montmorillonite and Bentonite

**Nuclear Energy and Safety Research Department (NES)
Laboratory for Waste Management (LES)**

M.A. Glaus, S. Frick, L.R. Van Loon

Paul Scherrer Institut
Forschungsstrasse 111
5232 Villigen PSI, Switzerland
Tel. +41 56 310 21 11
Fax +41 56 310 21 99
www.psi.ch

PSI Bericht Nr. 17-08
November 2017
ISSN 1019-0643

Table of Contents

Preface	I
Abstract	III
Zusammenfassung.....	V
Résumé	VII
摘要	IX
Table of Contents	XI
List of Tables.....	XIII
List of Figures	XV
List of Symbols and Abbreviations used in the present Report	XXXI
1 Introduction	1
1.1 Scope and aims of the report	1
1.2 Summary of the transport of charged species in charged argillaceous media focussing on smectite clay minerals.....	4
1.3 Report structure	9
2 Materials and Methods.....	11
2.1 Reagents, samples and analytical procedures	11
2.2 Diffusion experiments	11
2.3 Signal discrimination in multi-tracer experiments.....	15
2.3.1 Combination of γ - and β -emitters.....	15
2.3.2 β/β discrimination.....	16
2.4 Models for numerical simulations	17
2.5 Evaluation of best-fit parameter values and their uncertainties.....	18
3 Results for Diffusion in compacted Montmorillonite	21
3.1 Diffusion of HTO	21
3.2 Diffusion of cations: Bulk-dry densities of 1300 and 1600 kg m ⁻³	21
3.2.1 ²² Na ⁺ through- and out-diffusion (combined with HTO): 0.5 and 1.0 M NaClO ₄	21
3.2.2 ²² Na ⁺ through- and out-diffusion: 0.1 M NaClO ₄	24
3.2.3 ⁸⁵ Sr ²⁺ through- and out-diffusion: 0.5 and 1.0 M NaClO ₄	26
3.2.4 ⁸⁵ Sr ²⁺ through- and out-diffusion: 0.1 NaClO ₄	27
3.2.5 ¹³⁴ Cs ⁺ in-diffusion, various conditions.....	32
3.2.5.1 Concept and experimental procedures.....	32
3.2.5.2 Results: Diffusion of HTO.....	34
3.2.5.3 Results: Diffusion of ¹³⁴ Cs ⁺	34

3.3	Diffusion of $^{36}\text{Cl}^-$ (combined with HTO): Bulk-dry densities of 1300 and 1600 kg m^{-3}	39
3.4	Diffusion of cations: Bulk-dry density of 1900 kg m^{-3}	40
3.4.1	Diffusion of $^{85}\text{Sr}^{2+}$	40
3.4.2	Through-diffusion of $^{134}\text{Cs}^+$	44
3.5	Diffusion of anions: Bulk-dry density of 1900 kg m^{-3}	46
3.5.1	Through- and out-diffusion of $^{35}\text{SO}_4^{2-}$	46
3.5.2	Through- and out-diffusion of $^{75}\text{SeO}_4^{2-}$	47
4	Results for Diffusion in compacted Bentonite	49
4.1	Diffusion of HTO	49
4.2	Diffusion of $^{22}\text{Na}^+$	50
4.3	Diffusion of $^{85}\text{Sr}^{2+}$	51
4.4	Diffusion of $^{134}\text{Cs}^+$	53
4.5	Diffusion of $^{36}\text{Cl}^-$	55
4.6	Diffusion of $^{35}\text{SO}_4^{2-}$	57
5	Accessory Information	59
5.1	Profiles of water and anion contents in compacted montmorillonite	59
5.2	Re-saturation of compacted clay samples	60
5.3	Hydraulic tests	62
5.4	Batch sorption tests of $^{85}\text{Sr}^{2+}$ on dispersed Na- <i>mom</i>	63
5.5	Sorption measurements of $^{134}\text{Cs}^+$ on compacted Na- <i>mom</i> samples	64
6	Discussion	67
6.1	The "Gimmi – Kosakowski" scheme	67
6.2	Cation data	69
6.3	Anion data	72
6.4	Comparison of sorption results in compacted versus dispersed clay samples	74
6.5	Conclusions and implications for the selection of diffusion coefficients for radionuclides in performance assessment	75
6.6	General conclusions on 'surface-diffusion effects'	79
7	References	81
A	Appendix: Experimental data for Montmorillonite	A-1
A.1	Diffusion of HTO	A-1
A.2	Combined through-diffusion of $^{22}\text{Na}^+$ and HTO (0.5 and 1.0 M NaClO_4)	A-8
A.3	Combined out-diffusion of $^{22}\text{Na}^+$ and HTO (0.5 and 1.0 M NaClO_4)	A-16
A.4	Combined through- and out-diffusion of $^{22}\text{Na}^+$ and HTO (0.1 M NaClO_4)	A-24
A.5	Through- and out-diffusion of $^{85}\text{Sr}^{2+}$ (0.5 and 1.0 M NaClO_4)	A-26
A.6	^{134}Cs in-diffusion	A-33
A.7	Combined diffusion of HTO and $^{36}\text{Cl}^-$ (0.5 and 1.0 M NaClO_4)	A-39
A.8	^{85}Sr special test	A-47
A.9	Diffusion tests of $^{134}\text{Cs}^+$ in thin clay samples	A-55

A.10	Through- and out-diffusion of $^{35}\text{SO}_4^{2-}$	A-61
A.11	Through- and out-diffusion of $^{75}\text{SeO}_4^{2-}$	A-65
B	Appendix: Experimental data for Bentonite	B-1
B.1	Diffusion of HTO	B-1
B.2	Through- and out-diffusion of $^{22}\text{Na}^+$	B-9
B.3	Through- and out-diffusion of $^{85}\text{Sr}^{2+}$	B-15
B.4	Through-diffusion of $^{134}\text{Cs}^+$	B-21
B.5	Through- and out-diffusion of $^{36}\text{Cl}^-$	B-24
B.6	Through- and out-diffusion of $^{35}\text{SO}_4^{2-}$	B-27

List of Tables

Tab. 3.1:	Results of through-diffusion experiments with HTO in compacted Na- <i>mom</i> ^a	21
Tab. 3.2:	Results of through-diffusion experiments with $^{22}\text{Na}^+$ (and simultaneously measured HTO) in compacted Na- <i>mom</i> ^a	22
Tab. 3.3:	Results of the out-diffusion experiments with $^{22}\text{Na}^+$ (and simultaneously measured HTO) in compacted Na- <i>mom</i> ^a	23
Tab. 3.4:	Results of through-diffusion experiments with $^{22}\text{Na}^+$ (and simultaneously measured HTO) in compacted Na- <i>mom</i>	24
Tab. 3.5:	Results of the through- and out-diffusion experiments with $^{85}\text{Sr}^{2+}$ in compacted Na- <i>mom</i> ^a	27
Tab. 3.6:	Sensitivity analysis in the determination of D_e and α for a series of given D_f values in the diffusion experiments with $^{85}\text{Sr}^{2+}$ at a bulk-dry density of $\sim 1300 \text{ kg m}^{-3}$ and 0.1 M NaClO ₄ (cells 1058_5A, 1058_5B, 1058_7A, 1058_7B)	28
Tab. 3.7:	Sensitivity analysis in the determination of D_e and α for a series of given D_f values in the diffusion experiments with $^{85}\text{Sr}^{2+}$ at a bulk-dry density of $\sim 1300 \text{ kg m}^{-3}$ and 0.1 M NaClO ₄ (cells 1058_5A, 1058_5B, 1058_7A, 1058_7B)	30
Tab. 3.8:	Summary of the best-fit parameter values obtained from through-diffusion experiments of $^{85}\text{Sr}^{2+}$ in compacted Na- <i>mom</i> estimated from the range of possible values in a sensitivity analysis (<i>cf.</i> Tabs. 3.6 and 3.7)	30
Tab. 3.9:	Summary D_e values for HTO measured before in-diffusion of $^{134}\text{Cs}^+$	34
Tab. 3.10:	Summary D_e values for the in-diffusion of $^{134}\text{Cs}^+$	39
Tab. 3.11:	Results of the through-diffusion experiments with $^{36}\text{Cl}^-$ (and simultaneously measured HTO) in compacted Na- <i>mom</i> ^a	40
Tab. 3.12:	Results of combined through-diffusion experiments (labelled TON.1027) with $^{85}\text{Sr}^{2+}$ and HTO in compacted Na- <i>mom</i> in 1.0 M NaClO ₄	42

Tab. 3.13:	Results of through-diffusion experiments (labelled TON.1023) with $^{85}\text{Sr}^{2+}$ in compacted Na- <i>mom</i> at low external salinities ^a	44
Tab. 3.14:	Results of through-diffusion of $^{134}\text{Cs}^+$ combined with tracer profile analysis in compacted Na- <i>mom</i> (experiment TON.1034) ^a	45
Tab. 3.15:	Results of through- and out-diffusion experiments using $^{35}\text{SO}_4^{2-}$ in compacted Na- <i>mom</i> (experiment TON.1032) ^a	47
Tab. 3.16:	Results of through- and out-diffusion experiments using $^{75}\text{SeO}_4^{2-}$ in compacted Na- <i>mom</i> (experiment TON.1059) ^a	48
Tab. 4.1:	Results of through- and out-diffusion experiments of HTO in compacted Volclay ^a	50
Tab. 4.2:	Results of through-diffusion experiments combined with tracer profile analysis of $^{22}\text{Na}^+$ in compacted Volclay ^a	51
Tab. 4.3:	Results of through-diffusion experiments combined with tracer profile analysis or out-diffusion of $^{85}\text{Sr}^{2+}$ in compacted Volclay ^a	52
Tab. 4.4:	Results of through-diffusion experiments combined with tracer profile analysis of $^{134}\text{Cs}^+$ in compacted Volclay KWK ^a	54
Tab. 4.5:	Results of through- and out-diffusion experiments of $^{36}\text{Cl}^-$ in compacted Volclay KWK ^a	56
Tab. 4.6:	Results of through-diffusion of $^{35}\text{SO}_4^{2-}$ combined with tracer profile analysis in compacted Volclay KWK ^a	58
Tab. 5.1:	Average porosity values (\pm standard uncertainty) of Na- <i>mom</i> saturated with NaClO_4 calculated from profiling after through-diffusion.....	59
Tab. 5.2:	Average porosity values (\pm standard uncertainty) for the saturation of Na- <i>mom</i> with NaCl and NaClO_4 calculated from profiling after a saturation time of ~ 30 d.....	60
Tab. 5.3:	Determination of ρ_s in samples of compacted Volclay bentonite.....	62
Tab. 5.4:	Results of the sorption measurements of Cs^+ on compacted Na- <i>mom</i> samples (experiment TON.1068).....	66
Tab. 6.1:	Best-fit parameter values derived from regression (<i>cf.</i> Eq. (6.12)) of the data shown in Figs. 6.1 and 6.2. q is the minuend of the right-hand side of Eq. (6.12).....	75
Tab. 6.2:	Best-fit parameter values for the HTO diffusion data in Na- <i>mom</i> using the extended Archie relation (<i>cf.</i> Eq. (6.13)).....	78